

# CHS RAPID UP™

18-18-18 ■ With 1.75% Boron

## Guaranteed Analysis

Total Nitrogen (N) . . . . .	18.00%
2.30% Ammoniacal nitrogen	
4.10% Nitrate nitrogen	
11.60% Urea nitrogen	
Available Phosphate (P <sub>2</sub> O <sub>5</sub> ) . . . . .	18.00%
Soluble Potash (K <sub>2</sub> O) . . . . .	18.00%
Boron (B) . . . . .	1.75%
Copper (Cu) . . . . .	0.05%
0.05% Chelated copper	
Iron (Fe) . . . . .	0.10%
0.10% Chelated iron	
Manganese (Mn) . . . . .	0.05%
0.05% Chelated manganese	
Molybdenum (Mo) . . . . .	0.001%
Zinc (Zn) . . . . .	0.05%
0.05% Chelated zinc	

Derived from monoammonium phosphate, potassium nitrate, urea, monopotassium phosphate, sodium borate, boric acid, copper EDTA, iron EDTA, manganese EDTA, sodium molybdate, and zinc EDTA.

Chlorine (Cl) maximum . . . . . 0.025%  
Potential acidity is equivalent to 260lbs Calcium Carbonate (CaCO<sub>3</sub>) per ton.

## GENERAL INFORMATION

Applications of this product should be based on soil test and/or leaf analysis results. Consult your crop specialist for specific application rates. Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.aapfco.org/metals.html>

## DIRECTIONS FOR USE

Apply at the rate of 3 to 5 lbs per acre with sufficient water for good coverage as a supplement to normal soil fertility programs. This product contains 1.75% actual Boron and should only be used on crops requiring additional Boron. This product should not be used on Boron sensitive crops.

**Vegetable Crops:** As a foliar spray use 3 to 10 lbs. per acre. Applications may be made at 7-10 day interval depending on the amount of growth desired or when supplemental feeding is necessary. Depending on the crop, the first treatment should be made when the plants are 3-4 weeks old or when there is sufficient foliage for spraying. These rates may be applied to Vegetable Crops (tomatoes, peppers, cucumbers, beans, lettuce, celery, melons, squash, radishes, onions, broccoli, cabbage, spinach, cauliflower and potatoes.

**Raspberries And Blueberries:** Use at the rate of 3 to 5 lbs per acre early in the season and/or whenever it is necessary to improve cane vigor and growth. CAUTION: High rates of RAPID UP 18-18-18 applied late in the season may delay fruit color and maturity.

**Strawberries:** Use at the rate of 3 to 5 lbs per acre when the first fruit buds show in the crown. Two to three additional applications at 7-10 day intervals are beneficial.

**COMPATIBILITY:** RAPID UP 18-18-18 is compatible with most pesticides commonly used on vegetable, fruit and agronomic crops. CAUTION: Do not use with Dormant Oil, Lime Sulfur Spray Lime, or highly alkaline spray materials.

Add RAPID UP 18-18-18 to the spray tank first. Agitate in the tank is preferred to increase the speed of solubility. Where extremely cold water is used, allow a few extra minutes for the product to completely dissolve before spraying.

Use this product in accordance with good agronomic practices, which include utilizing proven spray equipment set for proper coverage. Do not make applications when temperatures are too hot. Applications should be made at temperature levels and when other environmental conditions in your area are such that your experience indicates the application will be compatible and will accomplish the desired result.

**WARNING:** This product contains boron (B); use on any crops other than those recommended may result in serious injury to the crops. The application of fertilizing materials containing molybdenum (Mo) may result in forage crops containing levels of molybdenum (Mo) which are toxic to ruminant animals. It also contains Copper, Iron, Manganese, and Zinc. Should be used as recommended.



**WARNING:** Harmful in contact with skin.  
Suspected of damaging fertility or the unborn child.

**Precautionary Statements:** Obtain special instructions before use. Do not handle until all safety precautions have been read and understood. Wear eye/face protection and protective gloves/clothing. Wash thoroughly after handling. **First Aid:** If on skin: Wash with plenty of water. If exposed or concerned, get medical advice/attention. Take off contaminated clothing and wash it before reuse. **Storage and Disposal:** Do not contaminate water, food or feed by storage or disposal. Keep out of reach of children. Store locked up. Dispose of contents/container in accordance with local regulations.

**WARNING:** Cancer and Reproductive Harm  
[www.P65Warningsca.gov](http://www.P65Warningsca.gov)

Conditions of Sale and Warranty Limitations: The Directions for Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of CHS Inc. or the seller. All such risks shall be assumed by the buyer. CHS Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use, subject to the inherent risks referred to above. CHS Inc. makes no other express or implied warranty of fitness or merchantability or any other express or implied warranty. In no cases shall CHS Inc. or the seller be liable for consequential, special or indirect damages resulting from the use or handling of this product. CHS Inc. and the seller offer this product, and the Buyer and the User accept it, subject to the foregoing conditions of Sale and Warranty, which may be varied only by agreement in writing signed by a duly authorized representative of CHS Inc.