

CHS RAPID UP™

20-20-20 Fruits/Vegetables Water Soluble Fertilizer

Guaranteed Analysis

Total Nitrogen (N)	20.00%
3.90% Ammoniacal Nitrogen	
5.80% Nitrate Nitrogen	
10.30% Urea Nitrogen	
Available Phosphate (P ₂ O ₅)	20.00%
Soluble Potash (K ₂ O)	20.00%
Boron (B)	0.020%
Copper (Cu)	0.050%
0.05% Chelated Copper	
Iron (Fe)	0.100%
0.10% Chelated Iron	
Manganese (Mn)	0.050%
0.05% Chelated Manganese	
Molybdenum (Mo)	0.0005%
Zinc (Zn)	0.050%
0.05% Chelated Zinc	

Derived from Monoammonium Phosphate, Potassium Nitrate, Urea, Sodium Borate, Copper EDTA, Iron EDTA, Manganese EDTA, Sodium Molybdate, and Zinc EDTA.
Chelating agent is Ethylenediaminetetraacetic Acid (EDTA).



WARNING KEEP OUT OF REACH OF CHILDREN

PRECAUTIONS

- Obtain special instructions before use.
- Do not handle until all safety precautions have been read and understood.
- Do not breathe dusts, mist or sprays. Wear appropriate NIOSH respirator.
- Avoid contact with skin. Wear long-sleeve shirt and disposable coveralls when contact potential is significant.
- Wash thoroughly after handling.
- Do not drink or smoke when using this product.
- Harmful if swallowed.
- May cause serious eye irritation.
- May cause skin irritation.

CAUTION: This product contains Boron, Copper, Iron, Manganese, Molybdenum, and Zinc. Should be used as recommended. Misuse may prove harmful. Do not use with highly alkaline spray materials, Bordeaux, or spray oils. Always add CHS Rapid Up™ 20-20-20 to spray mixture as the first ingredient.

NET WEIGHT 25 lbs./11.35 kg

RESPONSE

- IF IN EYES: Flush with water for 15 minutes. Call a doctor for treatment if irritation persists.
- IF ON SKIN: Wash thoroughly with soap and water. Launder clothing before reuse.
- IF INHALED: Remove affected person from source of exposure. Call 911 if breathing is difficult.
- IF INGESTED: Do not induce vomiting. Rinse mouth. Get medical attention.

EMERGENCY 800-210-9324

GENERAL INFORMATION

Applications of this product should be based on soil test and/or leaf analysis results. Consult your crop specialist for specific application rates. Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>.

DIRECTIONS FOR USE

GENERAL FOLIAR APPLICATION: Apply 2-10 lbs. of product per acre in a minimum 20 gallons of water.

AIRCRAFT APPLICATION: Apply a maximum of 4 lbs. of product in 2 gallons of water per acre.

VEGETABLE CROPS: As a foliar spray use 5-10 lbs. per acre. Applications may be made at 7-10 day interval depending on the amount of growth desired or when supplemental feeding is necessary. Depending on the crop, the first treatment should be made when the plants are 3-4 weeks old or when there is sufficient foliage for spraying. These rates may be applied to Vegetable Crops (tomatoes, peppers, cucumbers, beans, lettuce, celery, melons, squash, radishes, onions, broccoli, cabbage, spinach, cauliflower and potatoes.

FRUIT/VINE CROPS: As a foliar spray use 5-10 lbs. per acre. For dilute sprays use 2 lbs. per 100 gal. of water. Apply early in the season or when improved vigor is desired. 3-4 applications may be applied per season. Note: Do not use in late season sprays where fruit color and maturity are delayed by nitrogen applications. These rates may be applied to citrus (all kinds), apples, peaches, pears, avocados, prunes, apricots, cherries, nectarines, strawberries, hops and other vine crops. Apply early in the season during flush of new growth and well before fruit matures. Note: Use a minim of 100 gal. of water per acre for dilute sprays and not less than 5 gal. per acre for concentrated sprays.

GRAPES, RASPBERRIES, BLACKBERRIES AND BLUEBERRIES: Apply at the rate of 5 to 10 lbs. per acre early in the season or whenever it is necessary to improve vine or cane vigor and growth. For best results, repeat at pre-bloom and 3 weeks later. Use sufficient water for good coverage.

Conditions of Sale and Warranty Limitations: The Directions for Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of CHS Inc. or the seller. All such risks shall be assumed by the buyer. CHS Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use, subject to the inherent risks referred to above. CHS Inc. makes no other express or implied warranty of fitness or merchantability or any other express or implied warranty. In no cases shall CHS Inc. or the seller be liable for consequential, special or indirect damages resulting from the use or handling of this product. CHS Inc. and the seller offer this product, and the Buyer and the User accept it, subject to the foregoing conditions of Sale and Warranty, which may be varied only by agreement in writing signed by a duly authorized representative of CHS Inc.